



# On bipartite distance-regular graphs with a strongly closed subgraph of diameter three

Qian Kong, Kaishun Wang\*

School of Mathematical Sciences and Laboratory of Mathematics and Complex Systems, Beijing Normal University, Beijing, 100875, China

## ARTICLE INFO

### Article history:

Received 18 January 2010

Received in revised form 5 July 2010

Accepted 31 August 2010

Available online 22 September 2010

### Keywords:

Distance-regular graph

Strongly closed subgraph

## ABSTRACT

Let  $\Gamma$  denote a distance-regular graph with a strongly closed regular subgraph  $Y$ . Hosoya and Suzuki [R. Hosoya, H. Suzuki, Tight distance-regular graphs with respect to subsets, European J. Combin. 28 (2007) 61–74] showed an inequality for the second largest and least eigenvalues of  $\Gamma$  in the case  $Y$  is of diameter 2. In this paper, we study the case when  $\Gamma$  is bipartite and  $Y$  is of diameter 3, and obtain an inequality for the second largest eigenvalue of  $\Gamma$ . Moreover, we characterize the distance-regular graphs with a completely regular strongly closed subgraph  $H(3, 2)$ .

© 2010 Elsevier B.V. All rights reserved.

## 1. Introduction

Let  $\Gamma = (X, R)$  denote a simple connected graph with the vertex set  $X$  and the edge set  $R$ . For vertices  $x$  and  $y$ ,  $\partial(x, y)$  denotes the *distance* between  $x$  and  $y$ , i.e., the length of a shortest path connecting  $x$  and  $y$ . Let  $d := \max\{\partial(x, y) | x, y \in X\}$  denote the *diameter* of  $\Gamma$ . For  $x \in X$  and  $i \in \{0, 1, \dots, d\}$ , define

$$\Gamma_i(x) = \{y \in X | \partial(x, y) = i\}.$$

For two vertices  $x$  and  $y$  at distance  $i$ , define

$$C(x, y) = \Gamma_{i-1}(x) \cap \Gamma_1(y),$$

$$A(x, y) = \Gamma_i(x) \cap \Gamma_1(y),$$

and

$$B(x, y) = \Gamma_{i+1}(x) \cap \Gamma_1(y),$$

where  $\Gamma_{-1}(x) = \Gamma_{d+1}(x) = \emptyset$ .

A graph  $\Gamma$  is said to be *distance-regular* if the cardinalities  $c_i = |C(x, y)|$ ,  $a_i = |A(x, y)|$ , and  $b_i = |B(x, y)|$  are constants whenever  $\partial(x, y) = i$  ( $i = 0, 1, \dots, d$ ).

Let  $Y$  be a non-empty subset of  $X$ . When there is no danger of confusion, we also use  $Y$  to denote the induced subgraph on it.  $Y$  is called a *strongly closed* subgraph of  $\Gamma$  if  $C(x, y) \cup A(x, y) \subseteq Y$  whenever  $x, y \in Y$ . For  $x \in X$ , define

$$\partial(x, Y) = \min\{\partial(x, y) | y \in Y\}.$$

The numbers

$$\tau = \tau(Y) = \max\{\partial(x, Y) | x \in X\}, \quad w(Y) = \max\{\partial(x, y) | x, y \in Y\}$$

\* Corresponding author.

E-mail address: [wangks@bnu.edu.cn](mailto:wangks@bnu.edu.cn) (K. Wang).

are called the *covering radius*, the *width* of  $Y$  in  $\Gamma$ , respectively. For  $i \in \{0, 1, \dots, \tau\}$ , let

$$\Gamma_i(Y) = \{x \in X \mid \partial(x, Y) = i\}.$$

A subgraph  $Y$  is called *completely regular* if for any integer  $i$ ,

$$\gamma_i = |\Gamma_{i-1}(Y) \cap \Gamma_1(x)|, \quad \alpha_i = |\Gamma_i(Y) \cap \Gamma_1(x)|, \quad \beta_i = |\Gamma_{i+1}(Y) \cap \Gamma_1(x)|$$

are constants whenever  $x \in \Gamma_i(Y)$ .

Take  $\Omega$  to be a finite set of cardinality  $q \geq 2$ . The *Hamming graph*  $H(d, q)$  has vertex set  $\Omega^d$ , the Cartesian product of  $d$  copies of  $\Omega$ ; two vertices are adjacent whenever they differ in exactly one coordinate.

Let  $\Gamma$  denote a distance-regular graph with a regular subgraph  $Y$  of diameter 2. Hosoya and Suzuki [3] proved an inequality for the second largest and least eigenvalues of  $\Gamma$ . In this paper, we study the case when  $\Gamma$  is bipartite and  $Y$  is of diameter 3, and obtain the following results:

**Theorem 1.1.** *Let  $\Gamma = (X, R)$  be a bipartite distance-regular graph with a strongly closed regular subgraph  $Y$  of diameter 3. Then*

$$\theta_1 \leq \frac{b_2}{\sqrt{c_3 - c_2}}, \quad (1)$$

where  $\theta_1$  is the second largest eigenvalue of  $\Gamma$ .

**Proposition 1.2.** *Let  $\Gamma = (X, R)$  be a distance-regular graph with a strongly closed subgraph  $Y = H(3, 2)$ . If  $Y$  is completely regular with covering radius  $d - 3$ , then  $\Gamma$  is isomorphic to  $H(d, 2)$ .*

## 2. Preliminaries

In this section we recall some basic concepts about distance-regular graphs. Our notation and terminologies are standard. The reader is referred to [1,2] for more information.

In the rest of this paper, we always assume that  $\Gamma = (X, R)$  is a distance-regular graph of diameter  $d$  on  $n$  vertices. For each  $i$ , let  $A_i$  denote the matrix with  $(x, y)$ -entry

$$(A_i)_{xy} = \begin{cases} 1, & \text{if } \partial(x, y) = i, \\ 0, & \text{otherwise.} \end{cases}$$

We abbreviate  $A = A_1$ . The subalgebra  $\mathcal{M}$  of  $\text{Mat}_X(\mathbb{C})$  generated by  $A$  is called the *Bose–Mesner algebra* of  $\Gamma$ , and  $A_0, A_1, \dots, A_d$  form a basis of  $\mathcal{M}$ . It is well known that  $\mathcal{M}$  has a basis consisting of *primitive idempotents*  $E_0 = \frac{1}{n}J, E_1, \dots, E_d$ , where  $J$  is the all-one matrix of order  $n$ .

Let  $V = \mathbb{C}^X$  denote the vector space over the complex number field consisting of column vectors whose coordinates are indexed by  $X$  with complex entries. We endow  $V$  with the Hermitian inner product  $\langle \cdot, \cdot \rangle$  defined by

$$\langle \mathbf{u}, \mathbf{v} \rangle = {}^t \mathbf{u} \bar{\mathbf{v}},$$

where  ${}^t \mathbf{u}$  denotes the transpose of  $\mathbf{u}$ , and  $\bar{\mathbf{v}}$  is the complex conjugate of  $\mathbf{v}$ . We use the following abbreviation  $\|\mathbf{u}\|^2 = \langle \mathbf{u}, \mathbf{u} \rangle$ .

It is well-known that  $V$  has the following orthogonal decomposition

$$V = E_0 V \oplus E_1 V \oplus \dots \oplus E_d V,$$

where  $E_0 V = \langle \mathbf{1} \rangle$ ,  $\mathbf{1}$  is the all-one vector of size  $n$ . Let  $\hat{x}$  be the element of  $V$  with a 1 in the  $x$ -coordinate and 0 in all other coordinates. Moreover, if  $\Gamma$  is a bipartite distance regular graph with bipartitions  $P_1$  and  $P_2$ , then  $E_d V = \langle \hat{\mathbf{1}} \rangle$ , where  $\hat{\mathbf{1}} = \sum_{y \in P_1} \hat{y} - \sum_{z \in P_2} \hat{z}$ .

For each  $i$ , set

$$A_i = \sum_{j=0}^d p_i(j) E_j, \quad E_i = \frac{1}{n} \sum_{j=0}^d q_i(j) A_j.$$

Let  $m_i = q_i(0)$  and  $\theta_i = p_i(i)$ . Then  $m_i = \text{rank } E_i$ , and  $\theta_0, \theta_1, \dots, \theta_d$  are mutually distinct eigenvalues of  $\Gamma$ . By [1, p. 63],

$$\frac{p_i(j)}{k_i} = \frac{q_j(i)}{m_j},$$

where  $k_i = |\Gamma_i(x)|$ . We also write  $k_1 = k$ . For each eigenvalue  $\theta_j$ , let

$$\sigma_i = \sigma_i(\theta_j) = \frac{p_i(j)}{k_i} = \frac{q_j(i)}{m_j}. \quad (2)$$

We call the numbers  $\sigma_0, \sigma_1, \dots, \sigma_d$  the *cosine sequence* associated with  $\theta_j$ .

Let  $\theta$  be an eigenvalue of  $\Gamma$  and  $\sigma_0, \sigma_1, \dots, \sigma_d$  denote the corresponding cosine sequence. By [5, Lemma 2.1],

$$c_i\sigma_{i-1} + a_i\sigma_i + b_i\sigma_{i+1} = \theta\sigma_i,$$

and

$$\sigma_0 = 1, \quad \sigma_1 = \frac{\theta}{k}, \quad \sigma_2 = \frac{\theta^2 - a_1\theta - k}{kb_1}, \quad \sigma_3 = \frac{\theta^3 - (a_1 + a_2)\theta^2 + (a_1a_2 - k - c_2b_1)\theta + a_2k}{kb_1b_2}.$$

Substituting  $\sigma_1, \sigma_2, \sigma_3$  in (2) using the above equalities, we get

$$q_j(1) = \frac{m_j\theta}{k}, \quad (3)$$

$$q_j(2) = \frac{m_j(\theta^2 - a_1\theta - k)}{kb_1}, \quad (4)$$

$$q_j(3) = \frac{m_j(\theta^3 - (a_1 + a_2)\theta^2 + (a_1a_2 - k - c_2b_1)\theta + a_2k)}{kb_1b_2}. \quad (5)$$

For a non-empty set  $Y \subseteq X$  and  $x \in X$ , define  $E_0^* = E_0^*(Y)$  to be the diagonal matrix in  $\text{Mat}_X(\mathbb{C})$  with  $(x, x)$ -entry

$$(E_0^*)_{xx} = \begin{cases} 1, & \text{if } x \in Y, \\ 0, & \text{otherwise.} \end{cases}$$

A nonzero vector  $\mathbf{v} \in E_0^*V$  is said to be *tight* if

$$|\{i \mid i \in \{0, 1, \dots, d\}, E_i\mathbf{v} = \mathbf{0}\}| = w(Y).$$

For  $\mathbf{v} \in V \setminus \{\mathbf{0}\}$ , let

$$\eta^{(i)}(\mathbf{v}) = \frac{{}^t\mathbf{v}A_i\bar{\mathbf{v}}}{{}^t\mathbf{v}\mathbf{v}}, \quad \tilde{A}_i = E_0^*A_iE_0^*.$$

Pick  $\mathbf{v} \in E_0^*V$ . Then

$${}^t\mathbf{v}A_i\bar{\mathbf{v}} = {}^t\mathbf{v}E_0^*A_iE_0^*\bar{\mathbf{v}} = {}^t\mathbf{v}\tilde{A}_i\bar{\mathbf{v}}.$$

Moreover,  $\eta^{(i)}(\mathbf{v}) = 0$  for  $i > w(Y)$ . Furthermore, if  $\mathbf{v}$  is an eigenvector of  $\tilde{A}_i$ , then  $\eta^{(i)}(\mathbf{v})$  is the eigenvalue of  $\tilde{A}_i$  associated with the eigenvector  $\mathbf{v}$ .

### 3. Proof of main results

We begin with a useful lemma.

**Lemma 3.1.** Let  $\Gamma = (X, R)$  denote a bipartite distance-regular graph with a strongly closed regular subgraph  $Y$  of diameter 3, and let  $\mathbf{v} \in E_0^*V$  be a nonzero vector such that  $E_0\mathbf{v} = E_d\mathbf{v} = \mathbf{0}$ .

(i) For  $i \in \{0, 1, \dots, d\}$ ,

$$\frac{\|E_i\mathbf{v}\|^2}{\|\mathbf{v}\|^2} = -\frac{m_i}{kb_1b_2n}(\theta_i^2 - kb_1 - k)(\eta^{(1)}(\mathbf{v})\theta_i + b_2) \geq 0,$$

where  $k = \theta_0 > \theta_1 > \dots > \theta_d = -k$  are all distinct eigenvalues of  $\Gamma$ .

(ii) The following inequalities hold:

$$-\frac{b_2}{\theta_1} \leq -\sqrt{c_3 - c_2} \leq \eta^{(1)}(\mathbf{v}) \leq \sqrt{c_3 - c_2} \leq \frac{b_2}{\theta_1}.$$

(iii)  $\mathbf{v}$  is tight if and only if  $\eta^{(1)}(\mathbf{v}) = \pm\sqrt{c_3 - c_2} = \pm\frac{b_2}{\theta_1}$ .

**Proof.** (i) By [4, Lemma 8.2] and (3)–(5), we have

$$\begin{aligned} \frac{\|E_i\mathbf{v}\|^2}{\|\mathbf{v}\|^2} &= \frac{1}{n} \sum_{j=0}^{w(Y)} \eta^{(j)}(\mathbf{v})q_i(j) \\ &= \frac{m_i}{n} \left( 1 + \eta^{(1)}(\mathbf{v})\frac{\theta_i}{k} + \eta^{(2)}(\mathbf{v})\frac{\theta_i^2 - k}{kb_1} - (1 + \eta^{(1)}(\mathbf{v}) + \eta^{(2)}(\mathbf{v}))\frac{\theta_i^3 - (k + c_2b_1)\theta_i}{kb_1b_2} \right). \end{aligned}$$

Observe  $\eta^{(2)}(\mathbf{v}) = -1$  by  $E_d \mathbf{v} = \mathbf{0}$ , we find

$$\frac{\|E_i \mathbf{v}\|^2}{\|\mathbf{v}\|^2} = -\frac{m_i}{kb_1 b_2 n} (\theta_i^2 - kb_1 - k)(\eta^{(1)}(\mathbf{v})\theta_i + b_2).$$

(ii) For a vector  $\mathbf{u} \in V$ , let  $\mathbf{u}_Y$  be a vector with  $x$ th component

$$(\mathbf{u}_Y)_x = \begin{cases} \mathbf{u}_x, & \text{if } x \in Y, \\ 0, & \text{otherwise.} \end{cases}$$

Using  $E_0 \mathbf{v} = E_d \mathbf{v} = \mathbf{0}$ , we see  $\langle \mathbf{v}, \mathbf{1} \rangle = \langle \mathbf{v}, \tilde{\mathbf{1}} \rangle = 0$ . By  $\mathbf{v} \in E_0^* V$ ,  $\langle \mathbf{v}, \mathbf{1}_Y \rangle = \langle \mathbf{v}, \tilde{\mathbf{1}}_Y \rangle = 0$ . Suppose  $\eta$  is the second largest eigenvalue of  $\tilde{A}$ . Then

$$E_0^* V = \langle \mathbf{1}_Y \rangle \oplus \langle \tilde{\mathbf{1}}_Y \rangle \oplus W_\eta \oplus W_{-\eta}, \quad (6)$$

where  $W_{\pm\eta}$  is the eigenspace of  $\tilde{A}$  with corresponding eigenvalue  $\pm\eta$ . Then there exist  $\mathbf{v}_1, \mathbf{v}_2 \in E_0^* V$  such that

$$\mathbf{v} = \mathbf{v}_1 + \mathbf{v}_2,$$

where  $\tilde{A}\mathbf{v}_1 = \eta\mathbf{v}_1, \tilde{A}\mathbf{v}_2 = -\eta\mathbf{v}_2$ . Referring to [2, p. 432],  $\eta = \sqrt{c_3 - c_2}$ .

Note that

$${}^t \mathbf{v} \tilde{A} \tilde{\mathbf{v}} = {}^t \mathbf{v} \tilde{A} \tilde{\mathbf{v}} = \eta \|\mathbf{v}_1\|^2 - \eta \|\mathbf{v}_2\|^2,$$

and

$$\eta \|\mathbf{v}\|^2 \geq \eta \|\mathbf{v}_1\|^2 - \eta \|\mathbf{v}_2\|^2 \geq -\eta \|\mathbf{v}\|^2$$

reduce to

$$\eta \geq \eta^{(1)}(\mathbf{v}) \geq -\eta.$$

By (i), we obtain  $(\theta_i^2 - kb_1 - k)(\eta^{(1)}(\mathbf{v})\theta_i + b_2) \leq 0$ . If  $i = 0$  or  $d$ ,  $\theta_i^2 - kb_1 - k = 0$ . Then  $\theta_i^2 < k^2$  and  $\theta_i^2 - kb_1 - k < 0$  for  $1 \leq i \leq d-1$ . It follows that

$$\eta^{(1)}(\mathbf{v})\theta_i + b_2 \geq 0, \quad i = 1, \dots, d-1. \quad (7)$$

If  $\mathbf{v} = \mathbf{v}_1$  or  $\mathbf{v}_2$ , then  $\eta^{(1)}(\mathbf{v}) = \pm\eta$ , and

$$-\frac{b_2}{\theta_1} \leq \pm\eta \leq \frac{b_2}{\theta_1}.$$

(iii) Using  $E_0 \mathbf{v} = E_d \mathbf{v} = \mathbf{0}$ , the vector  $\mathbf{v}$  is tight if and only if there exists a unique  $i \in \{1, \dots, d-1\}$  such that  $E_i \mathbf{v} = \mathbf{0}$ . By (i) and  $\theta_i^2 - kb_1 - k < 0$ ,

$$\eta^{(1)}(\mathbf{v})\theta_i + b_2 = 0. \quad (8)$$

Referring to (7), the equality (8) holds if and only if  $i = d-1$  or  $i = 1$ , i.e.,

$$\eta^{(1)}(\mathbf{v}) = \pm\eta = \pm \frac{b_2}{\theta_1}. \quad \square$$

Next we start our proof of the main results.

**Proof of Theorem 1.1.** Referring to (6), we may pick linearly independent eigenvectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{|Y|-2}$  of  $\tilde{A}$  with corresponding eigenvalues  $\eta, -\eta$ . Moreover  $\mathbf{1}_Y, \tilde{\mathbf{1}}_Y$  are eigenvectors of  $\tilde{A}$  with corresponding eigenvalues  $c_3, -c_3$ , so  $\eta^{(1)}(\mathbf{1}_Y) = c_3$  and  $\eta^{(1)}(\tilde{\mathbf{1}}_Y) = -c_3$ . Since  $Y$  is bipartite distance-regular, we have

$$\text{tr}(\tilde{A}^2) = c_3|Y|, \quad |Y| = \frac{2(c_3^2 - c_3 + c_2)}{c_2},$$

which imply

$$\sum_{i=1}^{|Y|-2} (\eta^{(1)}(\mathbf{u}_i))^2 = c_3|Y| - 2c_3^2 = \frac{2c_3(c_3 - 1)(c_3 - c_2)}{c_2}. \quad (9)$$

Referring to Lemma 3.1(ii),

$$\sum_{j=1}^{|Y|-2} \left( \eta^{(1)}(\mathbf{u}_j) - \frac{b_2}{\theta_1} \right) \left( \eta^{(1)}(\mathbf{u}_j) + \frac{b_2}{\theta_1} \right) \leq 0. \quad (10)$$

Simplifying (10) using (9), (1) holds.  $\square$

Finally we shall characterize the distance-regular graphs with a completely regular strongly closed subgraph  $H(3, 2)$ .

Let  $v_0(t), v_1(t), \dots, v_d(t), v_{d+1}(t)$  denote the polynomials in  $\mathbb{R}[t]$  satisfying  $v_0(t) = 1$ , and for each  $i$ ,

$$tv_i(t) = b_{i-1}v_{i-1} + a_iv_i(t) + c_{i+1}v_{i+1}(t), \quad (11)$$

where  $b_{-1} = 0, c_{d+1} = 1$  and  $v_{-1}(t) = 0$ . Define

$$\rho_{1_Y}(t) = \frac{1}{n} \sum_{i=0}^d \eta^{(i)}(1_Y) \frac{v_i(t)}{k_i} \in \mathbb{R}[t]. \quad (12)$$

**Proof of Proposition 1.2.** By (11) we have

$$v_1(t) = t, \quad v_2(t) = \frac{1}{c_2}(t^2 - k), \quad v_3(t) = \frac{1}{c_2c_3}(t^3 - (k + b_1c_2)t).$$

Substituting  $v_1(t), v_2(t), v_3(t)$  in (12) using the above equalities, and then simplifying by  $k_{i-1}b_{i-1} = k_ic_i$ , we get

$$\begin{aligned} \rho_{1_Y}(t) &= \frac{1}{n} \sum_{i=0}^3 \eta^{(i)}(1_Y) \frac{v_i(t)}{k_i} \\ &= \frac{1}{n} \left( 1 + \eta^{(1)}(1_Y) \frac{t}{k_1} + \eta^{(2)}(1_Y) \frac{t^2 - k}{k_2c_2} + \eta^{(3)}(1_Y) \frac{t^3 - (k + b_1c_2)t}{k_3c_2c_3} \right) \\ &= \frac{1}{kb_1b_2n} (\eta^{(3)}(1_Y)t^3 + b_2\eta^{(2)}(1_Y)t^2 + (b_1b_2\eta^{(1)}(1_Y) - (k + b_1c_2)\eta^{(3)}(1_Y))t + kb_1b_2 - kb_2\eta^{(2)}(1_Y)). \end{aligned}$$

Observe

$$\eta^{(i)}(1_Y) = \frac{t 1_Y A_i \bar{1}_Y}{\|1_Y\|^2} = \frac{1}{|Y|} |\{(y_1, y_2) \in Y \times Y | \partial(y_1, y_2) = i\}|,$$

so  $\eta^{(1)}(1_Y) = \eta^{(2)}(1_Y) = 3, \eta^{(3)}(1_Y) = 1$  since  $Y$  is 3-cube. Moreover the fact that  $Y$  is strongly closed implies  $c_2 = 2, c_3 = 3, b_2 = k - 2$ . Hence,

$$\begin{aligned} \rho_{1_Y}(t) &= \frac{1}{kb_1b_2n} (t^3 + 3(k - 2)t^2 + (3k^2 - 12k + 8)t + k(k - 2)(k - 4)) \\ &= \frac{1}{kb_1b_2n} (t + k)(t + k - 2)(t + k - 4). \end{aligned} \quad (13)$$

By [4, Lemma 8.2] and (13),

$$\frac{\|E_i 1_Y\|^2}{\|1_Y\|^2} = \rho_{1_Y}(\theta_i) m_i = \frac{m_i}{kb_1b_2n} (\theta_i + k)(\theta_i + k - 2)(\theta_i + k - 4).$$

According to [3, Corollary 5.3],  $Y$  is completely regular with covering radius  $d - 3$  if and only if  $1_Y$  is tight, so  $-k, 2 - k$  and  $4 - k$  are the eigenvalues of  $\Gamma$ . In particular,  $\Gamma$  is a bipartite graph, and  $k - 2 = b_1 - 1$  is an eigenvalue of  $\Gamma$ . By the inequality in Theorem 1.1,  $\theta_1 \leq k - 2$ , so  $\theta_1 = k - 2$ . By [2, Theorem 4.4.11]  $\Gamma$  is isomorphic to  $H(d, 2)$ .  $\square$

**Remark.**  $H(d, 2)$ , Doubled Grassmann graph ( $q = 2$  or  $3$ ) with diameter 5 and doubled Odd graph on  $2m + 1$  points with  $m \geq 2$  satisfy the equality in Theorem 1.1.

## Acknowledgements

The authors would like to thank Professor Hiroshi Suzuki for helpful conversations and suggestions while preparing this paper. They wish to express their gratitude to the referees whose detailed reports have greatly improved the presentation of this paper. This research is supported by NCET-08-0052, NSF of China (10871027), and the Fundamental Research Funds for the Central Universities of China.

## References

- [1] E. Bannai, T. Ito, Algebraic Combinatorics I: Association Schemes, Benjamin/Cummings, California, 1984.
- [2] A.E. Brouwer, A.M. Cohen, A. Neumaier, Distance-Regular Graphs, Springer-Verlag, Berlin, Heidelberg, 1989.
- [3] R. Hosoya, H. Suzuki, Tight distance-regular graphs with respect to subsets, European J. Combin. 28 (2007) 61–74.
- [4] H. Suzuki, The Terwilliger algebra associated with a set of vertices in a distance-regular graph, J. Algebraic Combin. 22 (2005) 5–38.
- [5] H. Suzuki, On strongly closed subgraphs with diameter two and the  $Q$ -polynomial property, European J. Combin. 28 (2007) 167–185.